

Invented for life



BOSCH

Vivalytic SARS-CoV-2 DT, Flu A/B, RSV

Qualitative real-time PCR test for the detection of SARS-CoV-2, Flu A/B and Respiratory Syncytial Virus (RSV)

Vivalytic from Bosch –
The universal diagnostic platform.
Universal, intuitive, fast.



vivalytic

Vivalytic SARS-CoV-2 DT, Flu A/B, RSV

Qualitative real-time PCR test for the detection of SARS-CoV-2, Flu A/B and Respiratory Syncytial Virus (RSV)

The high burden caused each year by Covid-19, influenza and RSV infections shows how important it is to identify the different viruses quickly in order to provide the patient with targeted treatment.


Patients infected with SARS-CoV-2, influenza A (Flu A), influenza B (Flu B) and/or respiratory syncytial virus (RSV) have overlapping symptoms, but the approaches to patient management of infections caused by the viruses are different. The Vivalytic SARS-CoV-2 DT, Flu A/B, RSV test detects and differentiates between these viral nucleic acids, aiding in the diagnosis and thereby enabling pathogen-specific management of patients.

The SARS coronavirus 2 (SARS-CoV-2) is a respiratory pathogen which is responsible for the novel disease COVID-19. With over 10 million cases and 500,000 deaths reported as of end of June 2020, COVID-19 is challenging health care systems and society all over the world. It is important not only to have effective infection-preventing

measures in place, but also to trace and break the chains of infection through extensive testing.

Influenza, also called flu, is a contagious respiratory disease caused by influenza viruses. An influenza infection causes mild to severe illness and can be lethal. In Europe, seasonal influenza causes between 4 and 50 million symptomatic cases each year and the death toll associated with influenza is estimated to be between 15,000 and 70,000 every influenza season, in terms of excess deaths.¹

The respiratory syncytial virus (RSV) is a pathogen for acute diseases of the upper and lower respiratory tract. Especially infants and young children are at risk of developing a severe RSV infection resulting in hospitalization. Since clinical symptoms of RSV are nonspecific and can overlap with other viral respiratory infections such as influenza and SARS-CoV-2, differentiation of pathogens with other viral respiratory infections is needed.



Sample material

Nasopharyngeal, oropharyngeal or saliva swab samples


Volume
300 µl

Detectable pathogens


- SARS-CoV-2 virus
- Influenza A and Influenza B virus
- Human Respiratory Syncytial Virus (RSV)

Detection method
Real-time PCR test


Vivalytic workflow: Only a few steps from sample to result




① Scan sample code.



② Scan cartridge code.



③ Insert the sample and close cartridge lid.



④ Insert cartridge. The test starts automatically.

Our biocontent cooperation partner is:



Subject to technical modifications. Printed in Germany. Any colour deviations are due to restrictions in the printing process. Printed on 100% chlorine-free bleached cellulose, produced in an environmentally friendly process. Not all products are available in all regions. Ask your local sales representative for availability in specific markets. For in vitro diagnostic use only.

Bosch Healthcare Solutions GmbH
Stuttgarter Str. 130, 71332 Waiblingen, Germany

¹ European Centre for Disease Prevention and Control (ECDC)

The availability of the tests can differ regionally.

www.bosch-vivalytic.com

Art. No.: MDQ-01.733-01-LoC-009_Rev3.0